



STD	Symptoms	How you get it	Treatment	Outcome if not treated
<p>Chlamydia</p>	<p>Symptoms usually appear within 1 to 3 weeks after contact.</p> <p>For WOMEN:</p> <ul style="list-style-type: none"> - abnormal vaginal or rectal (if you have anal sex) discharge - burning sensation when urinating - abdominal or low back pain - nausea - fever - pain during sexual intercourse - bleeding between menstrual cycles <p>For MEN:</p> <ul style="list-style-type: none"> - discharge from the penis. - burning sensation around the opening of the penis when urinating. - pain and/or swelling in the testicles. 	<p>It is spread during vaginal, anal, or oral sex with someone infected with Chlamydia.</p>	<p>Chlamydia can be completely curable with antibiotics.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Reproductive organs can be severely and irreversibly damaged. - Women may no longer be able to have children. - Women may give Chlamydia to her baby during childbirth.
<p>Gonorrhea (a.k.a. – Clap, drip)</p>	<p>Symptoms usually appear within 2 to 21 days after contact.</p> <ul style="list-style-type: none"> - Thick yellow, gray, or green discharge from the vagina or penis. - Burning or pain when urinating or having a bowel movement. - Need to urinate more often. <p>For WOMEN:</p> <ul style="list-style-type: none"> - Abnormal periods or bleeding between periods. - Cramps and pain in the lower abdomen (belly). <p>For MEN:</p> <ul style="list-style-type: none"> - Swollen or tender testicles. 	<p>It is spread during vaginal, anal, or oral sex with someone infected with Gonorrhea.</p>	<p>Gonorrhea can be completely curable with antibiotics.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Cause infertility. - Arthritis. - Rash or skin diseases. - Heart trouble. - Kidney damage. - Blindness. - Women may give Gonorrhea to her baby during childbirth.
<p>Syphilis</p>	<p>1st Stage: Symptoms appear 1 to 12 weeks after contact.</p> <ul style="list-style-type: none"> - Sores on the mouth or sex organs <p>2nd Stage: Symptoms appear 1 week to 6 months after contact.</p> <ul style="list-style-type: none"> - Rash anywhere on the body. - Flu-like symptoms 	<p>It is spread during vaginal, anal, or oral sex with someone infected with Syphilis.</p>	<p>Syphilis can be completely curable with antibiotics. However, any damage done is not reversible.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Damage to internal organs including the brain, heart, blood vessels, liver, bones, and joints. - Women may give Syphilis to her baby during childbirth. - Blindness or death.

This brochure is not intended as a substitute for your health professional's opinion or care.

This publication, printed by DocuTech of the Oklahoma State Department of Health, is issued by the Oklahoma State Department of Health, as authorized by James M. Crutcher, MD, MPH, Commissioner of Health and State Health Officer. 50,000 copies have been printed at a cost of \$2,500.00. Copies have been deposited with the Publications Clearinghouse of the Oklahoma Department of Libraries.



STD	Symptoms	How you get it	Treatment	Outcome if not treated
<p>Herpes</p>	<p>Some people never have symptoms.</p> <p>Symptoms usually appear within 1 to 30 days or longer after contact.</p> <ul style="list-style-type: none"> - Flu-like symptoms. - Small, painful blisters on the sex organs or mouth. - Itching or burning before the blisters appear. - Blisters last 1 to 3 weeks. - Blisters may come and go, but you still have herpes. 	<p>It is spread during vaginal, anal, or oral sex or by genital touching with someone who has herpes.</p>	<p>There is currently no cure. There is medication that is available to relieve the pain of the blisters and prolong the time between outbreaks.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Women may give Herpes to her baby during childbirth.
<p>Genital Warts (HPV)</p>	<p>Symptoms usually appear within 1 to 8 months after contact.</p> <p>Small, bumpy warts on the penis, vagina, or anus.</p> <p>Itching or burning around the sex organs or anus.</p>	<p>It is spread during vaginal, anal, or oral sex or by genital touching with someone who has genital warts.</p>	<p>There is currently no cure. The warts may be removed by freezing or laser therapy.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Women may give Genital Warts (HPV) to her baby during childbirth.
<p>HIV/AIDS</p>	<p>Symptoms can appear anywhere from a few months to a few years after contact.</p> <ul style="list-style-type: none"> - Unexplained weight loss - Flu-like symptoms that don't go away. - Diarrhea. - White spots in the mouth. - In women, yeast infections that don't go away. 	<ul style="list-style-type: none"> - Spread during vaginal, anal, or oral sex with someone that has HIV. - Spread by sharing needles to inject drugs, or for any other reason. - Spread by contact with infected blood. 	<p>There is currently no cure. There is medication that postpones and/or treats specific symptoms.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Illness. - Women may give HIV/AIDS to her baby during childbirth. - Death.
<p>Hepatitis B</p>	<p>Symptoms usually appear within 1 to 9 months after contact. Although, some don't have any symptoms.</p> <ul style="list-style-type: none"> - Flu-like symptoms. - Tiredness. - Jaundice (yellow skin). - Dark urine (pee), light-colored bowel movements. 	<ul style="list-style-type: none"> - Spread during vaginal, anal, or oral sex with someone that has Hepatitis B. - Spread by sharing needles to inject drugs, or for any other reason. - Spread by contact with infected blood. 	<p>There is no cure. However, there is a vaccine for Hepatitis B.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Permanent liver damage or liver cancer. - Premature birth. - Women may give Hepatitis B to her baby during childbirth - Death.
<p>Trichomoniasis (a.k.a. Trich)</p>	<p>Many people never have symptoms.</p> <p>Symptoms can appear within 3 to 14 days after contact.</p> <p>In WOMEN:</p> <ul style="list-style-type: none"> - Itching, burning, or irritation in the vagina. - Yellowish, greenish, or gray discharge from the vagina. <p>In MEN:</p> <ul style="list-style-type: none"> - Watery, white drip from the penis. - Burning or pain when urinating (peeing). - Need to urinate (pee) more often. 	<p>Spread during vaginal sex.</p>	<p>Trichomoniasis can be completely curable with antibiotics.</p>	<p>If it is not treated, any of the following can result:</p> <ul style="list-style-type: none"> - Give the disease to your sex partner(s). - Men can get prostate infections. - Symptoms will continue.