

PROPER USE OF COMBINATION VACCINES

Brand Name	Antigens	Licensed for:	Schedule	Minimum & Maximum Ages	Special Notes
COMVAX®	Hep B - Hib	All 3 doses of the Hib & Hep B series	2, 4, & 12-15 mos.	6 weeks through 4 years of age	Must not be used for infants younger than 6 weeks of age
Pediarix®	DTaP- Hep B - IPV	1 st 3 doses of DTaP & IPV series	2, 4, & 6 mos.	6 weeks through 6 years of age	Cannot be used for the birth dose of Hep B or 4th dose of DTaP series
Pentacel®	DTaP-IPV/Hib	1 st 4 doses of DTaP, IPV and Hib series	2, 4, 6, & 12-18 mos.	6 weeks through 4 years of age	Do not administer to children age 5 years or older
ProQuad®	MMRV	1 st & 2 nd doses of measles, mumps, rubella, and varicella	12-15 mos. & 4-6 yrs.	12 months of age through 12 years of age	Do not administer to children, adolescents or adults age 13 years or older
TriHIBit®	DTaP- Hib	4 th dose of DTaP series and 4 th dose of Hib series	12-18 mos.	12 months through 4 years of age	Must not be used for the primary series at 2, 4, or 6 months of age
Kinrix®	DTaP-IPV	5 th dose of DTaP series and 4 th dose of polio series	4-6 yrs.	4 through 6 years of age	Not for use for the primary series at 2, 4, or 6 months of age or the 4th DTaP at 12 – 18 months of age