

## OK by One Schedule & Beyond Using Combination Vaccines

### OK by One Schedule Using Pediarix, ActHIB or PedvaxHIB, Hiberix, or TriHIBit & Kinrix

| Vaccine                        | Birth | 2 months            | 4 months            | 6 months   | 12 months | At least 6 months later | 4-6 years                                      |  |        |
|--------------------------------|-------|---------------------|---------------------|--|-----------|-------------------------|--|--|--------|
| Hep B                          | Hep B | Pediarix            | Pediarix            | Pediarix   |           |                         |  |  |        |
| IPV                            |       |                     |                     |  |           |                         |  |  | Kinrix |
| DTaP                           |       |                     |                     |  |           |                         | TriHIBit OR DTaP & PedvaxHib OR DTaP & Hiberix |  |        |
| Hib                            |       | ActHIB or PedvaxHib | ActHIB or PedvaxHib | ActHib or None needed if 1 <sup>st</sup> 2 doses were both PedvaxHib |           |                         |  |  |        |
| PCV                            |       | PCV                 | PCV                 | PCV  | PCV       |                         |  |  |        |
| RV (PO)                        |       | RotaTeq or Rotarix  | RotaTeq or Rotarix  | RotaTeq or None needed if 1 <sup>st</sup> 2 doses were both Rotarix  |           |                         |  |  |        |
| MMR                            |       |                     |                     |  | MMR       |                         | MMR  |  |        |
| Varicella                      |       |                     |                     |  | Var       |                         | Var  |  |        |
| Hep A                          |       |                     |                     |  | Hep A     | Hep A                   |  |  |        |
| Total # of injections at Visit | 1     | 3                   | 3                   | 2 or 3   | 5 or 6    | 1                       | 3  |  |        |
| Influenza                      |       |                     |                     | 2 doses 4 weeks apart if receiving for the first time                |           |                         |  |  |        |

Total number of injections will vary based on the type of Hib used and the availability and use of ProQuad (combined MMR and Varicella).

Administer influenza vaccine to all children 6 months through 18 years of age. Give two doses to all children 6 months through 8 years who are receiving influenza vaccine for the first time or who failed to receive the 2<sup>nd</sup> dose in the preceding year. Give doses at least 4 weeks apart.

### Ok by One Schedule Using Pentacel & ActHib or Hiberix

| Vaccine                        | Birth | 2 months           | 4 months           | 6 months  | 12 months         | At least 6 months later | 4-6 years            |
|--------------------------------|-------|--------------------|--------------------|---|-------------------|-------------------------|----------------------|
| Hep B                          | Hep B | Hep B              |                    | Hep B   |                   |                         |                      |
| DTaP                           |       | Pentacel           | Pentacel           | Pentacel  | DTaP              |                         | DTaP & IPV or Kinrix |
| IPV                            |       |                    |                    |   |                   |                         |                      |
| Hib                            |       |                    |                    |   | ActHIB OR Hiberix |                         |                      |
| PCV                            |       | PCV                | PCV                | PCV   | PCV               |                         |                      |
| RV (PO)                        |       | RotaTeq or Rotarix | RotaTeq or Rotarix | RotaTeq or None needed if 1 <sup>st</sup> 2 doses were both Rotarix |                   |                         |                      |
| MMR                            |       |                    |                    |   | MMR               |                         | MMR                  |
| Varicella                      |       |                    |                    |   | Var               |                         | Var                  |
| Hep A                          |       |                    |                    |   | Hep A             | Hep A                   |                      |
| Total # of injections at Visit | 1     | 3                  | 2                  | 3   | 5                 | 1                       | 4                    |
| Influenza                      |       |                    |                    | 2 doses 4 weeks apart if receiving for the first time               |                   |                         |                      |

Total number of injections will vary based on availability and use of ProQuad (combined MMR and Varicella).

Administer influenza vaccine to all children 6 months through 18 years of age. Give two doses to all children 6 months through 8 years who are receiving influenza vaccine for the first time or who failed to receive the 2<sup>nd</sup> dose in the preceding year. Give doses at least 4 weeks apart.

OK by One Schedule Using COMVAX & Kinrix

| Vaccine                        | Birth | 2 months           | 4 months           | 6 months  | 12 months | At least 6 months later | 4-6 years            |
|--------------------------------|-------|--------------------|--------------------|---|-----------|-------------------------|----------------------|
| Hep B                          | Hep B | COMVAX             | COMVAX             | None needed at this visit   | COMVAX    |                         |                      |
| Hib                            |       |                    |                    |   |           |                         |                      |
| DTaP                           |       | DTaP               | DTaP               | DTaP  | DTaP      |                         | Kinrix or DTaP & IPV |
| IPV                            |       | IPV                | IPV                | IPV   |           |                         |                      |
| PCV                            |       | PCV                | PCV                | PCV   | PCV       |                         |                      |
| RV (PO)                        |       | RotaTeq or Rotarix | RotaTeq or Rotarix | RotaTeq or None needed if 1 <sup>st</sup> 2 doses were both Rotarix |           |                         |                      |
| MMR                            |       |                    |                    |   | MMR       |                         | MMR                  |
| Varicella                      |       |                    |                    |   | Var       |                         | Var                  |
| Hep A                          |       |                    |                    |   | Hep A     | Hep A                   |                      |
| Total # of Injections at Visit | 1     | 4                  | 4                  | 3   | 6         | 1                       | 3                    |
| Influenza                      |       |                    |                    | 2 doses 4 weeks apart if receiving for the first time               |           |                         |                      |

Total number of injections will vary based on availability and use of ProQuad (combined MMR and Varicella).

Administer influenza vaccine to all children 6 months through 18 years of age. Give two doses to all children 6 months through 8 years who are receiving influenza vaccine for the first time or who failed to receive the 2<sup>nd</sup> dose in the preceding year. Give doses at least 4 weeks apart.

OK by One Schedule Without Comvax, Pentacel, Pediarix or Kinrix

| Vaccine                        | Birth | 2 months            | 4 months            | 6 months   | 12 months                      | At least 6 months later | 4-6 years |
|--------------------------------|-------|---------------------|---------------------|--|--------------------------------|-------------------------|-----------|
| Hep B                          | Hep B | Hep B               |                     | Hep B  |                                |                         |           |
| DTaP                           |       | DTaP                | DTaP                | DTaP   | DTaP                           |                         | DTaP      |
| Hib                            |       | ActHIB or PedvaxHib | ActHIB or PedvaxHib | ActHib or None needed if 1 <sup>st</sup> 2 doses were both PedvaxHib | ActHIB OR PedvaxHib OR Hiberix |                         |           |
| IPV                            |       | IPV                 | IPV                 | IPV  |                                |                         | IPV       |
| PCV                            |       | PCV                 | PCV                 | PCV  | PCV                            |                         |           |
| RV (PO)                        |       | RotaTeq or Rotarix  | RotaTeq or Rotarix  | RotaTeq or None needed if 1 <sup>st</sup> 2 doses were both Rotarix  |                                |                         |           |
| MMR                            |       |                     |                     |  | MMR                            |                         | MMR       |
| Varicella                      |       |                     |                     |  | Var                            |                         | Var       |
| Hep A                          |       |                     |                     |  | Hep A                          | Hep A                   |           |
| Total # of Injections at Visit | 1     | 5                   | 4                   | 4 or 5   | 5                              | 1                       | 4         |
| Influenza                      |       |                     |                     | 2 doses 4 weeks apart if receiving for the first time                |                                |                         |           |

Total number of injections may vary based on the type of Hib vaccine used and the availability and use of ProQuad (combined MMR and Varicella).

Administer influenza vaccine to all children 6 months through 18 years of age. Give two doses to all children 6 months through 8 years who are receiving influenza vaccine for the first time or who failed to receive the 2<sup>nd</sup> dose in the preceding year. Give doses at least 4 weeks apart.

## Our Clinic Schedule (Fill in the blanks with the vaccines your clinic prefers or routinely uses.)

| Vaccine               | Birth | 2 months | 4 months | 6 months  | 12 months | At least 6 months later | 4-6 years |
|-----------------------|-------|----------|----------|---|-----------|-------------------------|-----------|
| Hep B                 | HepB  |          |          |   |           |                         |           |
| DTaP                  |       |          |          |   |           |                         |           |
| Hib                   |       |          |          |   |           |                         |           |
| IPV                   |       |          |          |   |           |                         |           |
| PCV                   |       | PCV      | PCV      | PCV   | PCV       |                         |           |
| RV (PO)               |       |          |          |   |           |                         |           |
| MMR                   |       |          |          |   | MMR       |                         | MMR       |
| Varicella             |       |          |          |   | Var       |                         | Var       |
| Hep A                 |       |          |          |   | Hep A     | Hep A                   |           |
| Total # of Injections | 1     |          |          |   |           |                         |           |
| Influenza             |       |          |          | 2 doses 4 weeks apart if receiving for the first time |           |                         |           |

### Notes:

- Four doses of Hepatitis B vaccine may be administered when a birth dose is given.
- TriHIBit cannot be used in the primary series at 2, 4, or 6 months of age. It may be used as the Hib booster (final) dose following a primary series of single antigen Hib vaccine or Hib vaccine in combination with other vaccines - meaning TriHIBit can be used if the child is aged  $\geq 12$  months **and** has received at least one prior dose of Hib vaccine  $\geq 2$  mos. earlier, **and** TriHIBit will be the last dose in the Hib series.
- Hiberix should not be used for the primary series of Hib vaccination at 2, 4 and 6 months of age. Hiberix can be used as the booster or final dose of the Hib series following a primary series of single antigen Hib vaccine or Hib vaccine in combination with other vaccines - meaning Hiberix can be used if the child is aged  $\geq 12$  months **and** has received at least one prior dose of Hib vaccine  $\geq 2$  mos. earlier, **and** Hiberix will be the last dose in the Hib series.
- Total number of injections will vary based on availability and use of ProQuad (combined MMR and Varicella) and type of Hib vaccine used.
- Administer influenza vaccine to all children 6 months through 18 years of age. Give two doses to all children 6 months through 8 years who are receiving influenza vaccine for the first time or who failed to receive the 2<sup>nd</sup> dose in the preceding year. Give doses at least 4 weeks apart.