



Muskogee County

Unintentional Poisoning Facts

State Facts:

- From 2007-2013, more than 4,600 Oklahomans died of an unintentional poisoning (UP).
- More Oklahoma adults age 25-64 die of UP than motor vehicle crashes or suicide.
- Opioid prescription painkillers are the most common drugs involved in UP deaths.
- Oklahoma had the sixth highest poisoning death rate in the U.S.

What is a Poisoning?

A poisoning is the ingestion, inhalation, absorption, or contact with a substance resulting in a toxic effect or bodily harm. An unintentional poisoning occurs when a person does not intend to hurt themselves or someone else. The person may intentionally be exposed to a substance (e.g., a person ingests a medication at higher than prescribed levels) but does not intend to harm themselves.

In the late 1990s the most common cause of overdose deaths became prescription drugs.

Nearly four out of five unintentional poisoning deaths in Oklahoma involve at least one prescription drug.

Number of UP deaths in Muskogee County by type of substance, 2007-2013:

All unintentional poisonings	119
Prescription drugs	98
Prescription painkillers (opioids)	88
Illicit drugs	29
Alcohol	10

Most common substances* in Muskogee County overdose deaths:

Oxycodone	Morphine
Alprazolam	Methamphetamine
Hydrocodone	Fentanyl
Methadone	Cocaine
Diazepam	Alcohol

Injury Prevention Service

(405) 271-3430
<http://poison.health.ok.gov>



Muskogee County Data (2007-2013)

- From 2007-2013, there were 119 unintentional poisoning deaths in Muskogee County.
- Muskogee County had the 7th highest unintentional poisoning death rate in the state.
- Nearly three-fourths of these deaths involved a prescription painkiller.
- Males were more likely to die of an unintentional poisoning than females.
- Adults age 35-54 had the highest rate of unintentional poisoning death.
- Adults age 45-54 were six times as likely to die of an unintentional poisoning compared to teens and young adults age 15-24.
- Four out of ten people who died of an unintentional poisoning had a history of mental health problems.
- Nearly six out of ten people who died had a history of substance abuse.
- Six out of ten deaths occurred at a home or apartment, while one in three occurred at a hospital.

*Top three substances in bold.

Prevention

- Take medications as prescribed, and never more than the recommended dosage
- Never share or sell prescription drugs
- Properly dispose of unused, unneeded, or expired drugs
- Keep all medication in a safe place to avoid theft
- Call 211 for help finding treatment referrals

Signs/Symptoms of an Overdose

- Won't awaken when aroused
- Bluish purple skin tones for lighter skinned people and grayish or ashen tones for darker skinned people
- Slow, shallow, erratic, or absent breathing
- Snore-like gurgling or choking sounds
- Elevated body temperature
- Vomiting
- Irrational behavior or confusion

Emergency: Call 911
immediately if you suspect
someone is overdosing.




Common Generic/ Brand Name Prescription Drugs

<i>Generic</i>	<i>Brand Name</i>
<i>Alprazolam</i>	<i>Xanax</i>
<i>Carisoprodol</i>	<i>Soma</i>
<i>Cyclobenzaprine</i>	<i>Flexeril, Fexmid</i>
<i>Diazepam</i>	<i>Valium</i>
<i>Fentanyl</i>	<i>Duragesic patch</i>
<i>Hydrocodone</i>	<i>Lortab, Norco, Vicodin</i>
<i>Methadone</i>	<i>Diskets, Methadose</i>
<i>Morphine</i>	<i>MS Contin, Kadian, Avinza</i>
<i>Oxycodone</i>	<i>OxyContin, Percocet, Endocet</i>
<i>Tramadol</i>	<i>Ultram, Rybix, Ryzolt</i>
<i>Zolpidem</i>	<i>Ambien, Intermezzo</i>

(SIGNATURE)

REFILL 0 | 2 3 4 5 PRN

Unintentional Poisoning Death Rates by County of Residence¹, Oklahoma, 2007-2013



¹County of residence was unknown for 39 persons.

Source: OSDH, Injury Prevention Service, Unintentional Poisonings Database
 (Abstracted from Medical Examiner reports)