


HIV/AIDS CASES AMONG BLACKS



OVERVIEW

In this fact sheet, race was reclassified and these numbers will not match the number of HIV/AIDS cases among Blacks previously reported. The race category calculation in other fact sheets follows the CDC recommendation, where persons reporting Hispanic ethnicity are classified as Hispanic and persons reporting two or more races are classified as Multi Race. For the purpose of this analysis, any person reporting one of their races as Black were reclassified as Black, instead of Multi Race, and persons reporting race as Black and ethnicity as Hispanic were reclassified as Black, instead of Hispanic.

Blacks accounted for 23.1% (2,256) of the 9,783 cumulative HIV/AIDS cases diagnosed in Oklahoma at the end of 2014. The breakdown of the cumulative cases diagnosed among Blacks was 57.1% (1,289) diagnosed with AIDS and 42.9% (967) diagnosed with only HIV. Of the cumulative cases, 34.1% (769) were known to be deceased. Blacks accounted for 26.5% (1,487) of the 5,610 cases living with HIV/AIDS in Oklahoma at the end of 2014. Of the living cases among Blacks, 43.5% (647) were diagnosed with AIDS and 56.5% (840) were diagnosed with HIV. Of the 311 newly diagnosed HIV cases in 2014, Blacks accounted for 29.3% (91). Of the newly diagnosed cases among Blacks, 80.2% (73) were diagnosed with HIV and 19.8% (18) were diagnosed with HIV and AIDS in 2014.

BY SEX


Males accounted for 79.1% (1,785) of the cumulative cases diagnosed among Blacks in Oklahoma and females accounted for the remaining 20.9% (471) of cases. Among

Blacks living with HIV/AIDS in Oklahoma, males accounted for 77.1% (1,147) of the cases and females accounted for 22.9% (340). Among the newly diagnosed HIV cases among Blacks, males accounted 72.5% (66) of the cases and females accounted for 27.5% (25).

BY AGE

By age at time of diagnosis, the 20-29 years age group (610; 41.0%) accounted for the largest number of living HIV/AIDS cases among Blacks, followed by 30-39 years (437; 29.4%) and 40-49 years (229; 15.4%). Teenagers (13-19 years) accounted for 6.6% (98) of the cases. By current age group, the 40-49 years age group (376; 25.3%) has the highest burden of living HIV/AIDS cases among Blacks in Oklahoma, followed very closely by 50-59 years of age (352; 23.7%), 30-39 years of age (333; 22.4%), and 20-29 years of age (284; 19.1%).

Living HIV/AIDS Cases by Current Age Group Among Blacks, Oklahoma 2014



For More Information Contact:

HIV/STD Service
Surveillance and
Analysis

1000 NE 10th Street,
MS 0308
Oklahoma City, OK 73117

405.271.4636

hivstd.health.ok.gov



Oklahoma State
Department of Health


HIV/AIDS CASES AMONG BLACKS

Males accounted for **77.1%** of the living cases among Blacks.

30-39 years age group accounted for **1 in 5** newly diagnosed HIV cases.

Just over 45% (41; 45.1%) of the newly diagnosed HIV cases among Blacks were 20-29 years of age at diagnosis. The 30-39 years age group (18; 19.8%) accounted for nearly 20% of the cases. Teenagers accounted for 12% of the newly diagnosed cases among Blacks (11; 12.1%). The 40-49 years (9; 9.9%) and 50-59 years (9; 9.9%) age groups each accounted for nearly 10% of the cases.

Newly Diagnosed HIV Cases by Age Group Among Blacks, Oklahoma 2014



BY MODE OF TRANSMISSION.....


Of the 1,487 living HIV/AIDS cases among Blacks:

- 48.0% (714) were MSM¹,
- 16.5% (246) were heterosexual sex,
- 5.9% (87) were IDU,
- 4.8% (72) were MSM and IDU,
- 0.8% (12) were perinatal,
- 0.5% (8) were blood recipients, and
- 23.4% (348) were no reported risk or no identified risk.

Among the newly diagnosed HIV cases among Blacks, MSM accounted for the majority of the cases (50; 54.9%). Heterosexual sex, the second largest risk group, accounted for 15.4% (14) of the newly diagnosed cases. Other risk categories accounted for

4.4% (4) of the cases. Risk was not reported or identified for 25.3% (23) of the newly diagnosed cases.

Living HIV/AIDS Cases by Risk Among Blacks, Oklahoma 2014



BY GEOGRAPHY.....

Four counties accounted for over 80% (1,208; 81.2%) of the living HIV/AIDS cases diagnosed among Blacks in Oklahoma. Of these four counties, Oklahoma County accounted for the highest number of cases (640; 43.0%), followed by Tulsa County (366; 24.6%), Cleveland County (105; 7.1%), and Comanche County (97; 6.5%).

Four counties combined to account for over 75% (69; 75.8%) of the newly diagnosed HIV cases among Blacks. Oklahoma County (33; 36.3%) accounted for the highest number of cases, followed by Tulsa County (26; 28.6%). Canadian County and Cleveland County each accounted for 5.5% (5) of the cases.

¹Men who have sex with men

²Injection drug use